

Scattering from an AdS_3 bubble and an exact AdS_3

H. W. Lee and Y. S. Myung

Department of Physics, Inje University, Kimhae 621-749, Korea

Abstract

We investigate the close relationship between the potential and absorption cross section for test fields in an AdS_3 bubble (a 5D black hole) and an exact AdS_3 . There are two solutions in type IIB string theory: an AdS_3 bubble corresponds to the dilatonic solution, while an exact AdS_3 is the non-dilatonic solution. In order to obtain the cross section for an AdS_3 bubble, we introduce the {out}-state scattering picture with the AdS_3 -AFS matching procedure. For an exact AdS_3 , one considers the {in}-state scattering picture with the AdS_3 - AdS_3 matching. Here the non-normalizable modes are crucially taken into account for the matching procedure. It turns out that the cross sections for the test fields in an AdS_3 bubble take the same forms as those in an exact AdS_3 . This suggests that in the dilute gas and the low energy limits, the S-matrix for an AdS_3 bubble can be derived from an exact AdS_3 space.

I. INTRODUCTION

There has been a great progress in string theory of the D1-D5 brane system with momentum modes along the string direction(S^1). This gives us a 5D black hole(M_5) with three charges (Q_1, Q_5, Q_n) . The first progress was achieved in the Bekenstein-Hawking entropy [1]. Apart from the success of counting the microstates of a 5D black hole through D-brane physics, a dynamical consideration becomes an important issue [2–5]. This is so because the semiclassical absorption cross section (greybody factor) for a test field arises as a consequence of its potential barrier surrounding the horizon. That is, this is an effect of spacetime curvature. More precisely, it is worth noting that a semiclassical absorption cross section can be derived from a solution to the differential equation of a test field(most often, $\bar{\nabla}^2\phi = 0$) on the supergravity side.

The AdS/CFT correspondence states that string theory in the AdS is dual to a conformal field theory(CFT) defined on its remote boundary of AdS [6]. The semiclassical limit of spacetime physics is related to the large N limit of the dual CFT. A 5D black hole($M_5 \times S^1 \times T^4$) becomes $AdS_3 \times S^3 \times T^4$ near horizon but with an asymptotically flat space(AFS) [7]. Recently this is called an AdS_3 bubble in AFS and corresponds to the dilatonic solution [8]. In this case the matching procedure is crucial for obtaining an absorption coefficient and here we need to match an AdS_3 bubble to AFS.

On the other hand, one obtains $AdS_3 \times S^3 \times T^4$ as the other solution to the type IIB string theory. This is an exact AdS_3 with asymptotically AdS_3 and corresponds to the non-dilatonic solution [9]. Further it includes the BTZ black hole [10]. We point out that the near horizon limit of a 5D black hole yields the BTZ black hole, whereas this solution accomodates to the BTZ black hole spacetime as a whole.

It was understood that the greybody factor calculation makes sense when one finds an asymptotically flat region as in Sec.II [6]. Hence it may not be possible in an AdS_3 because there is no asymptotic state corresponding to particle at infinity of AdS_3 . However, the authors in [9] calculated the greybody factor for a free scalar and the dilaton both

in $M_5 \times S^1 \times T^4$ (an AdS_3 bubble) and $AdS_3 \times S^3 \times T^4$ (an exact AdS_3) within the type IIB supergravity. Here in the exact AdS_3 calculation we choose the non-normalizable modes to obtain the greybody factor. This corresponds to the AdS_3 - AdS_3 matching procedure. This expression denotes shorthand for a certain choice of boundary conditions where non-normalizable modes inject flux into AdS_3 . It turns out that two results of a free scalar are exactly the same. And the results for the dilaton are the same upto a factor 3. These show a close relationship between two approaches. More recently, vacuum correlators of the dual CFT_4 were expressed as truncated n-point functions for the non-normalizable modes in AdS_5 . One can interpret this result as an S matrix of an exact AdS_5 arising from a limit of scattering from an AdS_5 bubble [8]. This supports that our calculation based on the non-normalizable modes is correct.

In this paper we will show that the S-matrix of an AdS_3 bubble can be derived partly from an exact AdS_3 space. This is one of the current issues in the AdS/CFT correspondence. For this purpose, we investigate the close relationship between the potential and absorption cross section in an AdS_3 bubble and an exact AdS_3 . Comparing an AdS_3 bubble with an exact AdS_3 leads to an assumption that the potential of an exact AdS_3 is the left-hand side of an AdS_3 bubble. For this study, we introduce the {in} and {out}-state pictures for an AdS_3 bubble. For the exact AdS_3 study, one needs the {in}-state as well as non-normalizable modes. Further, we introduce the test fields for scattering analysis. These are in an AdS_3 bubble : a free scalar(ϕ) which, in the decoupling limit, relates to an (1,1) operator $\mathcal{O}$ in the holographically dual theory; two fixed scalars(ν, λ) to (2,2), (3,1), and (1,3) operators; two intermediate scalars(η, ξ) to (1,2) and (2,1) operators. On the exact AdS_3 side, the test fields are a free scalar(ψ), the dilaton(Φ), an intermediate scalar(η), and the tachyon(T).

The organization of this work is as follows. Sec.II is devoted to analyzing the scattering from an AdS_3 bubble within a 5D black hole. This corresponds to a conventional scattering study. We study the scattering of the test fields in an exact AdS_3 in Sec. III. In this case we are careful for obtaining the absorption coefficient because we cannot define the asymptotically flat space. Finally we discuss our results in Sec. IV.

II. SCATTERING FROM AN ADS_3 BUBBLE IN A 5D BLACK HOLE

Initially we introduce all perturbing modes in a 5D black hole background. It is pointed out that in s-wave calculation the fixed scalars are physically propagating modes and other fields belong to be redundant modes [5]. Hence we choose two fixed scalars (ν, λ) and a free scalar (ϕ) as the relevant modes. We begin with the 5D black hole with three charges,

$$ds_{5\text{D}}^2 = -h f^{-2/3} dt^2 + f^{1/3} (h^{-1} dr^2 + r^2 d\Omega_3^2), \quad (1)$$

where

$$f = f_1 f_2 f_n = \left(1 + \frac{r_1^2}{r^2}\right) \left(1 + \frac{r_5^2}{r^2}\right) \left(1 + \frac{r_n^2}{r^2}\right), \quad h = \left(1 - \frac{r_0^2}{r^2}\right). \quad (2)$$

Here the radii are related to the boost parameters (α_i) and the charges (Q_i) as

$$r_i^2 = r_0^2 \sinh^2 \alpha_i = \sqrt{Q_i^2 + \frac{r_0^4}{4}} - \frac{r_0^2}{2}, \quad i = 1, 5, n. \quad (3)$$

Hence the D-brane black hole depends on the four parameters (r_1, r_5, r_n, r_0) . The event horizon (outer horizon) is clearly at $r = r_0$. When all three charges are nonzero, the surface $r = 0$ becomes a smooth inner horizon (Cauchy horizon). When at least one of the charges is zero, the surface $r = 0$ becomes singular. The extremal case corresponds to the limit of $r_0 \rightarrow 0$ with the boost parameters $\alpha_i \rightarrow \pm\infty$, keeping the charges (Q_i) fixed. We are interested in the limit of $r_0, r_n \ll r_1, r_5$, which is called the dilute gas limit. This is so because this limit corresponds to the near-outer horizon. Here we choose $Q_1 = r_1^2, Q_5 = r_5^2$, and $r_n = r_0 \sinh \alpha_n$ with a finite α_n . This corresponds to the near-extremal black hole and its thermodynamic quantities (energy, entropy, Hawking temperature) are given by

$$E_{\text{next}} = \frac{2\pi^2}{\kappa_5^2} \left[r_1^2 + r_5^2 + \frac{1}{2} r_0^2 \cosh 2\alpha_n \right], \quad (4)$$

$$S_{\text{next}} = \frac{4\pi^3 r_0}{\kappa_5^2} r_1 r_5 \cosh \alpha_n, \quad (5)$$

$$\frac{1}{T_{\text{H,next}}} = \frac{2\pi}{r_0} r_1 r_5 \cosh \alpha_n, \quad (6)$$

where κ_5^2 is the 5D gravitational constant. The above energy and entropy are those of a gas of massless 1D particles. In this case the temperatures for left and right moving string modes are given by

$$T_L = \frac{1}{2\pi} \left(\frac{r_0}{r_1 r_5} \right) e^{\alpha_n}, \quad T_R = \frac{1}{2\pi} \left(\frac{r_0}{r_1 r_5} \right) e^{-\alpha_n}. \quad (7)$$

This implies that the (left and right moving) momentum modes along the string direction are excited, while the excitations of D1-anti D1 and D5-anti D5-branes are suppressed. The Hawking temperature is given by their harmonic average

$$\frac{2}{T_H} = \frac{1}{T_L} + \frac{1}{T_R}. \quad (8)$$

A. Potential analysis

For a free scalar ($\phi = \phi(r)e^{i\omega t}Y_l(\theta_1, \theta_2, \theta_3)$), the linearized equation $\bar{\nabla}^2 \phi = 0$ in the background of (1) leads to [2,3,11]

$$\left[\left(h r^3 \partial_r \right)^2 + \omega^2 r^6 f - \frac{l(l+2)h}{r^2} \right] \phi = 0. \quad (9)$$

The s-wave ($l = 0$) linearized equation for the fixed scalars takes the form [4,5]

$$\left[\left(h r^3 \partial_r \right)^2 + \omega^2 r^6 f - \frac{8 h r^4 r_{\pm}^4}{(r^2 + r_{\pm}^2)^2} \left(1 + \frac{r_0^2}{r_{\pm}^2} \right) \right] \phi_{\pm} = 0, \quad (10)$$

where one gets ν for ϕ_+ and λ for ϕ_- . Here $r_{\pm}^2 = [r_1^2 + r_5^2 + r_n^2 \pm \sqrt{r_1^4 + r_5^4 + r_n^4 - r_1^2 r_5^2 - r_1^2 r_n^2 - r_5^2 r_n^2}]/3$. Considering $N = r^{-3/2} \tilde{N}$, for $N = \nu, \lambda, \phi$ and introducing a tortoise coordinate $r^* = \int (dr/h) = r + (r_0/2) \ln |(r - r_0)/(r + r_0)|$ [2], then the equation takes the form

$$\frac{d^2 \tilde{N}}{dr^{*2}} + (\omega^2 - \tilde{V}_N) \tilde{N} = 0. \quad (11)$$

Here we take $r_1 = r_5 = R$ and $r_0 = r_n$ for simplicity. In the dilute gas limit ($R \gg r_0$), $\tilde{V}_N(r)$ is given by

$$\tilde{V}_\nu(r) = -\omega^2(f-1) + h \left[\frac{3}{4r^2} \left(1 + \frac{3r_0^2}{r^2} \right) + \frac{8R^4}{r^2(r^2 + R^2)^2} \right], \quad (12)$$

$$\tilde{V}_\lambda(r) = -\omega^2(f-1) + h \left[\frac{3}{4r^2} \left(1 + \frac{3r_0^2}{r^2} \right) + \frac{8R^4}{r^2(3r^2 + R^2)^2} \right], \quad (13)$$

$$\tilde{V}_\phi(r) = -\omega^2(f-1) + h \left[\frac{3}{4r^2} \left(1 + \frac{3r_0^2}{r^2} \right) + \frac{l(l+2)}{r^2} \right], \quad (14)$$

where

$$f-1 = \frac{r_0^2 + 2R^2}{r^2} + \frac{(2r_0^2 + R^2)R^2}{r^4} + \frac{r_0^2 R^4}{r^6}. \quad (15)$$




FIG. 1. The graph of $(f-1)$ in 5D black hole with $r_0 = 0.01, R = 0.3$. A peak appears at outer horizon($r = r_0$).

We note that $\tilde{V}_N$ depends on two parameters (r_0, R) as well as the energy (ω) . As (11) stands, it is far from the Schrödinger-type equation. The ω -dependence is a matter of peculiar interest to us compared with the Schwarzschild black hole potentials (V_{RW}, V_Z, V_ψ) [12]. This makes the interpretation of $\tilde{V}_N$ as a potential difficult. As is shown in Fig.1, this

arises because $(f-1)$ is very large as 10^6 for $r_0 = 0.01, R = 0.3$ in the near-horizon. In order for $\tilde{V}_N$ to be a potential, it is necessary to take the low energy limit of $\omega \rightarrow 0$. It is suitable to be 10^{-3} . And $\omega^2(f-1)$ is of order $\mathcal{O}(1)$ and thus it can be ignored in comparison to the remaining ones. Now we can define a potential $V_N = \tilde{V}_N + \omega^2(f-1)$. Hence, in the low energy limit ($\omega \rightarrow 0$), Eq.(11) becomes as the Schrödinger-type equation. Further the last terms in (12)-(13) are important to compare each other. After the partial fraction, these lead to

$$\frac{8R^4}{r^2(r^2 + R^2)^2} = \frac{8}{r^2} - \frac{8}{r^2 + R^2} - \frac{8R^2}{(r^2 + R^2)^2}, \quad (16)$$

$$\frac{8R^4}{r^2(3r^2 + R^2)^2} = \frac{8}{r^2} - \frac{24}{3r^2 + R^2} - \frac{24R^2}{(3r^2 + R^2)^2}. \quad (17)$$

The last term of a free scalar in (14) with $l = 2$ keeps the first terms in (16)-(17) only. One finds immediately the sequence

$$V_{\phi_0} \ll V_\lambda \leq V_\nu \leq V_{\phi_2}. \quad (18)$$




FIG. 2. Four potential graphs ($V_\phi^{l=0}, V_\nu, V_\lambda, V_\phi^{l=2}$) for an AdS_3 bubble in a 5D black hole with $r_0 = 0.01, R = 0.3$.

Here ϕ_0 denotes the s-wave($l = 0$) free scalar and ϕ_2 the free one with $l = 2$. This is also observed from the graphs of potential in Fig.2 with $r_0 = 0.01, R = 0.3$. Because the shape of their potentials takes nearly the same form ($V_\lambda \simeq V_\nu \simeq V_\phi^{l=2}$), these give us nearly the same reflection coefficient $\mathcal{R} = |R|^2$ and absorption one $\mathcal{A} = |A|^2$. For example, in the low energy limit of $\omega \rightarrow 0$, λ, ν, ϕ_2 take nearly the zero-absorption cross section. Furthermore all potentials go to zero, as r approaches infinity. This implies the existence of the asymptotic states outside an AdS_3 bubble. The size of an AdS_3 bubble is from $r_0 = 0.01$ to $R = 0.3$.

B. Scattering from an AdS_3 bubble

We are interested in the scattering of the test fields off $V_N(r^*)$. It is well known that the scattering analysis is usually done by choosing a coordinate such as $-\infty \leq r^* \leq \infty$. It is always possible to visualize the black hole as presenting an effective potential barrier (or well) to an incoming test wave. One expects that some of the incident wave will be irreversibly absorbed by the black hole, while the remaining fraction will be scattered back to the infinity. In this scattering we can calculate the reflection and transmission(absorption) coefficients [13]. As is shown in Fig.3, we note that all potential barriers $V_N(r^*)$ take nearly the symmetric forms around $r^* = 0$ when they are rewritten by a tortoise coordinate r^* [14]. Also they are localized at $r^* = 0$. Since $V_N(r^*) \rightarrow 0$ as $r^* \rightarrow \pm\infty$, one finds

$$\frac{d^2 \tilde{N}_{\pm\infty}}{dr^{*2}} + \omega^2 \tilde{N}_{\pm\infty} = 0. \quad (19)$$

Asymptotically($r^* \rightarrow \infty$) the solution is given by

$$N_{+\infty}^{\text{out}} = e^{i\omega r^*} + R_N^{\text{out}}(\omega) e^{-i\omega r^*}. \quad (20)$$

Considering the time-part of $e^{i\omega t}$, the first is an incoming wave($\leftarrow$) and the last is an outgoing wave($\rightarrow$). Near the horizon it is purely incoming($\leftarrow$) as

$$N_{-\infty}^{\text{out}} = A_N^{\text{out}}(\omega) e^{i\omega r^*}. \quad (21)$$

We call this type of solution as $\{\text{out}\}_N$ and the corresponding vacuum state is defined as $b_i|0\rangle_{\text{out}} = 0$.




FIG. 3. Four potential graphs ($V_\phi^{l=0}, V_\nu, V_\lambda, V_\phi^{l=2}$) as functions of r^* for an AdS_3 bubble in a 5D black hole with $r_0 = 0.01, R = 0.3$.

In order to study the Hawking radiation, we introduce another boundary condition. Asymptotically the wave is purely outgoing($\rightarrow$)

$$N_{+\infty}^{\text{in}} = A_N^{\text{in}}(\omega)e^{-i\omega r^*} \quad (22)$$

but near the horizon it has both outgoing($\rightarrow$) and incoming($\leftarrow$) parts as

$$N_{-\infty}^{\text{in}} = e^{-i\omega r^*} + R_N^{\text{in}}(\omega)e^{i\omega r^*}. \quad (23)$$

We call this type of solution as $\{\text{in}\}_N$ and its vacuum state is defined as $a_i|0\rangle_{\text{in}} = 0$. The vacuum states $|0\rangle_{\text{out}}$ and $|0\rangle_{\text{in}}$ form two different bases of which any state can be expanded in terms of the other. These are two different Fock space vacuum states and $\{\text{out}\}_N$ and $\{\text{in}\}_N$ are related to each other by the Bogoliubov transformation,

$$b_i = \sum_j (\alpha_{ij}^* a_j - \beta_{ij}^* a_j^\dagger), \quad (24)$$

$$b_i^\dagger = \sum_j (\alpha_{ij} a_j^\dagger - \beta_{ij} a_j). \quad (25)$$

The computation of Hawking shows in a semi-classical approximation that the thermal radiation from the black hole with temperature T_H is given by [15]

$$\langle N_\omega \rangle = {}_{\text{in}} \langle 0 | b_i^\dagger b_i | 0 \rangle_{\text{in}} = \sum_k |\beta_{ik}|^2 = \frac{\sigma_{5\text{D}}^N}{e^{\omega/T_H} - 1}, \quad (26)$$

with an absorption cross section $\sigma_{5\text{D}}^N = |A_N^{\text{out}}|^2 \times 4\pi/\omega^3 = \mathcal{A}_N^{\text{out}} 4\pi/\omega^3$. Note that if $\sigma_{5\text{D}}^N$ is a constant, $\langle N_\omega \rangle$ is the same as that of a black body. Typically, $\sigma_{5\text{D}}^N$ is not constant but varies. The deviations from the black body spectrum have earned it the name “greybody factor”. Here we define the S-matrix from $\{\text{in}\}_N$ and $\{\text{out}\}_N$ as [14]

$$S(\omega) = \begin{pmatrix} A_N^{\text{out}}(\omega) & A_N^{\text{in}}(\omega) \\ R_N^{\text{out}}(\omega) & R_N^{\text{in}}(\omega) \end{pmatrix}. \quad (27)$$

Here an incident wave($e^{i\omega r^*}$) of unit amplitude from $r^* = +\infty$ gives rise to $A_N^{\text{out}}(\omega)$ and $R_N^{\text{out}}(\omega)$. On the other hand, an incident wave($e^{-i\omega r^*}$) of unit amplitude from $r^* = -\infty$ gives rise to $A_N^{\text{in}}(\omega)$ and $R_N^{\text{in}}(\omega)$. The relation between these is given by

$$A_N^{\text{out}}(\omega) = A_N^{\text{in}}(\omega) = A_N(\omega), \quad (28)$$

$$\frac{R_N^{\text{out}}(\omega)}{A_N(\omega)} = -\frac{R_N^{\text{in}}(-\omega)}{A_N(-\omega)}, \quad \frac{R_N^{\text{out}}(-\omega)}{A_N(-\omega)} = -\frac{R_N^{\text{in}}(\omega)}{A_N(\omega)}, \quad (29)$$

$$A_N^*(\omega) = A_N(-\omega), \quad R_N^{\text{out}*}(\omega) = R_N^{\text{out}}(-\omega), \quad R_N^{\text{in}*}(-\omega) = R_N^{\text{in}}(\omega). \quad (30)$$

The above relations establish the symmetry and unitarity of the S-matrix in the AdS₃ bubble scattering. Actually $A_N(\omega)$ can be calculated from the backscattering of an incident wave N off the potential $V_N(r^*)$. It is not easy to find out the absorption amplitude A_N directly in the complicated potentials such as $V_N(r^*)$. In this case one use the flux $\mathcal{F}$ of the incoming wave to obtain the absorption coefficient

$$\mathcal{A}_N^{\text{out}} = \frac{\mathcal{F}(-\infty)}{\mathcal{F}(\infty)}, \quad (31)$$

where $\mathcal{F}(-\infty)(\mathcal{F}(\infty))$ are the fluxes at the horizon(infinity). In this way we can calculate the semiclassical absorption cross section.

C. Exact analysis of near-horizon (AdS₃ bubble)

First let us consider a free scalar. Using $z = h$, the wave equation (9) can be rewritten as

$$z(1-z)\frac{d^2\phi}{dz^2} + (1-z)\frac{d\phi}{dz} + \left\{-C + \frac{Q}{z} + \frac{E}{1-z}\right\}\phi = 0, \quad (32)$$

where

$$C = \left(\frac{\omega r_1 r_5 r_n}{2r_0^2}\right)^2 = \frac{\omega^2}{64\pi^2} \left(\frac{1}{T_L} - \frac{1}{T_R}\right)^2, \quad (33)$$

$$E = -\frac{l(l+2)}{4} + \frac{\omega^2(r_1^2 + r_5^2 + r_n^2)}{4} \simeq -\frac{l(l+2)}{4} \quad (34)$$

$$Q = \left(\frac{\omega}{4\pi T_H}\right)^2 \left[\left(1 + \frac{r_0^2}{r_1^2} + \frac{r_0^2}{r_5^2}\right) + 4\pi^2 r_n^2 T_H^2 \right] \simeq \left(\frac{\omega}{4\pi T_H}\right)^2. \quad (35)$$

Here $\simeq$ means both the dilute gas limit ($r_0, r_n \ll r_1, r_5$) and low energy limit ($\omega \rightarrow 0$). Then one has $\omega r_0, \omega r_n \ll \omega r_1, \omega r_5 < 1$. In order to compare (32) with the hypergeometric equation, one has to transform it into the pole-free equation. With an unknown constant A , we find the ingoing mode at the horizon

$$\phi^n = A z^{-i\sqrt{Q}} (1-z)^{(1-\nu)/2} F(a, b, c; z), \quad (36)$$

where

$$\nu = \sqrt{(l+1)^2 - \omega^2(r_1^2 + r_5^2 + r_n^2)} \simeq l+1, \quad (37)$$

$$a = \frac{1-\nu}{2} - i\sqrt{Q} + i\sqrt{C} \simeq -\frac{l}{2} - i\frac{\omega}{4\pi T_R}, \quad (38)$$

$$b = \frac{1-\nu}{2} - i\sqrt{Q} - i\sqrt{C} \simeq -\frac{l}{2} - i\frac{\omega}{4\pi T_L}, \quad (39)$$

$$c = 1 - 2i\sqrt{Q} \simeq 1 - i\frac{\omega}{2\pi T_H}. \quad (40)$$

The large r -behavior ($z \rightarrow 1$) of ϕ^n can be obtained from the ($z \rightarrow 1-z$) transformation rule for the hypergeometric functions as

$$\phi^{n \rightarrow f} = \frac{A\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} u^{\nu-1} + \frac{A\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} u^{-(\nu+1)}. \quad (41)$$

For the fixed scalars (ν, λ), considering both (16), (17) and the near-horizon condition of $r \simeq r_0 \ll r_1, r_5$, one finds that (10) leads to (9) with $l = 2$. Thus one can obtain their near-horizon behaviors from ϕ_2 .

D. Asymptotic states

Let us first consider a free scalar. In the far region, we introduce $\phi = \check{\phi}/r$ and $u = \omega r$ and then (9) leads to

$$\frac{d^2 \check{\phi}}{du^2} + \frac{1}{u} \frac{d\check{\phi}}{du} + \left[1 - \frac{\nu^2}{u^2}\right] \check{\phi} = 0. \quad (42)$$

The solution is given by the Bessel function when ν is not an integer

$$\phi^f = \left[\alpha \frac{J_\nu(u)}{u} + \beta \frac{J_{-\nu}(u)}{u} \right], \quad (43)$$

where α, β are unknown constants. From the large u -behavior ($r \rightarrow \infty, \omega r \gg 1$), one finds the asymptotic states

$$\phi_\infty^f = \sqrt{\frac{1}{2\pi}} \frac{e^{-iu}}{u^{3/2}} \left\{ \alpha e^{i(\nu+1/2)\pi/2} + \beta e^{-i(\nu+1/2)\pi/2} \right\} \quad (44)$$

and its incoming flux

$$\mathcal{F}(\infty) = -\frac{2}{\omega^2} \left| \alpha e^{i(\nu+1/2)\pi/2} + \beta e^{-i(\nu+1/2)\pi/2} \right|^2. \quad (45)$$

The small u -behavior ($\omega r < 1$) of ϕ^f is

$$\phi^{f \rightarrow \text{inter}} = \frac{1}{u} \left[\alpha \left(\frac{u}{2}\right)^\nu \frac{1}{\Gamma(\nu+1)} + \beta \left(\frac{u}{2}\right)^{-\nu} \frac{1}{\Gamma(-\nu+1)} \right]. \quad (46)$$

On the other hand, the asymptotic behavior of the s-wave fixed scalars ($\phi_\pm = \check{\phi}_\pm/r$) is governed by

$$\frac{d^2 \check{\phi}_\pm}{du^2} + \frac{1}{u} \frac{d\check{\phi}_\pm}{du} + \left[1 - \frac{\nu^2}{u^2}\right] \check{\phi}_\pm = 0 \quad (47)$$

with $\nu \simeq 1$. Its solution is given by

$$\phi_\pm^f \simeq \left[\alpha_\pm \frac{J_\nu(u)}{u} + \beta_\pm \frac{J_{-\nu}(u)}{u} \right]. \quad (48)$$

Here $\alpha_\pm, \beta_\pm$ are unknown constants.

E. AdS₃-AFS matching procedure and absorption cross section

Here we use the matching procedure between an AdS₃ bubble (near-horizon of a 5D black hole) and asymptotically flat space to obtain an absorption coefficient. First consider the matching of a free scalar. Here the matching point resides on $0 < r^* < \infty$ in Fig.3. In the intermediate zone ($u < 1$), from (41) and (46) one finds

$$\alpha = Au_0 \frac{2^\nu \Gamma(\nu+1) \Gamma(\nu) \Gamma(c) u_0^{-\nu}}{\Gamma(c-a) \Gamma(c-b)}, \beta = Au_0 \frac{\Gamma(-\nu+1) \Gamma(-\nu) \Gamma(c) u_0^\nu}{2^\nu \Gamma(a) \Gamma(b)}. \quad (49)$$

Since $u_0 = \omega r_0 \ll 1$, one finds $\alpha \gg \beta$. In this case we take an incoming flux effectively. Furthermore the incoming flux at the horizon is found as

$$\mathcal{F}(0) = -8\pi r_0^2 \sqrt{Q} |A|^2. \quad (50)$$

The absorption coefficient is given by

$$\mathcal{A}_\phi^{\text{out}} = \frac{\mathcal{F}(0)}{\mathcal{F}(\infty)} \simeq 4\pi u_0^2 \sqrt{Q} \left| \frac{A}{\alpha} \right|^2. \quad (51)$$

The absorption cross section takes the form [11,16]

$$\begin{aligned} \sigma_{5D}^\phi &= (l+1)^2 \frac{4\pi}{\omega^3} \mathcal{A}_\phi^{\text{out}} \\ &\simeq \frac{\mathcal{A}_H^{5D}}{[l!(l+1)!]^2} (l+1)^2 \left(\frac{\omega r_0}{2} \right)^{2l} \left| \frac{\Gamma(\frac{l+2}{2} - i\frac{\omega}{4\pi T_L}) \Gamma(\frac{l+2}{2} - i\frac{\omega}{4\pi T_R})}{\Gamma(1 - i\frac{\omega}{2\pi T_H})} \right|^2 \end{aligned} \quad (52)$$

with the area of horizon $\mathcal{A}_H^{5D} = 2\pi r_1 r_5 r_n$. We have, for even l

$$\begin{aligned} \sigma_{5D}^{\phi_l} &= (l+1)^2 \frac{\pi^3}{2^{4l}} \frac{(r_1 r_5)^{2l+2} \omega^{2l+1}}{[l!(l+1)!]^2} [\omega^2 + (2\pi T_L)^2 2^2] \cdots [\omega^2 + (2\pi T_L)^2 l^2] \times \\ &\quad [\omega^2 + (2\pi T_R)^2 2^2] \cdots [\omega^2 + (2\pi T_R)^2 l^2] \frac{e^{\omega/T_H} - 1}{(e^{\omega/2T_L} - 1)(e^{\omega/2T_L} - 1)}. \end{aligned} \quad (53)$$

Matching procedure for the s-wave $\phi_\pm$ is nearly the same as in a free scalar [5]. It leads to

$$\sigma_{5D}^{\phi_\pm} = \frac{\pi^3 r_1^6 r_5^6}{64 r_\pm^4} \omega (\omega^2 + 16\pi^2 T_L^2) (\omega^2 + 16\pi^2 T_R^2) \frac{e^{\omega/T_H} - 1}{(e^{\omega/2T_L} - 1)(e^{\omega/2T_L} - 1)}. \quad (54)$$

In the limit of $\omega \ll T_L, T_R, T_H$, the low-energy absorption cross sections are calculated as

$$\sigma_{5D}^{\phi_0} = \mathcal{A}_H^{5D}, \quad (55)$$

$$\sigma_{5D}^{\phi_2} = \frac{3}{16} \mathcal{A}_H^{5D} (\omega r_0)^4 = \frac{3}{4} (\omega R)^4 \left\{ \frac{\mathcal{A}_H^{5D}}{4} \left(\frac{r_0}{R} \right)^4 \right\}, \quad (56)$$

$$\sigma_{5D}^\nu = \frac{\mathcal{A}_H^{5D}}{4} \left(\frac{r_0}{R} \right)^4, \quad (57)$$

$$\sigma_{5D}^\lambda = 9 \frac{\mathcal{A}_H^{5D}}{4} \left(\frac{r_0}{R} \right)^4, \quad (58)$$

where we impose the relation $r_1 = r_5 = R, r_0 = r_n$. Here we find a sequence of cross section

$$\sigma_{5D}^{\phi_0} \gg \sigma_{5D}^\lambda \geq \sigma_{5D}^\nu \geq \sigma_{5D}^{\phi_2}. \quad (59)$$

This originates from the potential sequence in (18). It is consistent with our naive expectation that the absorption cross section increases, as the height of potential decreases. Here we wish to point out the difference between a free scalar and the fixed scalars. In the dilute gas limit ($R \gg r_0$) and the low energy limit ($\omega \rightarrow 0$), the s-wave cross section for a free scalar ($\sigma_{5D}^{\phi_0}$) goes to $\mathcal{A}_H^{5D}$ [3], while the s-wave cross sections for fixed scalars (ν, λ) including ϕ_2 approach zero [4]. Also this can be confirmed from Fig.2.

III. SCATTERING FROM AN EXACT AdS_3 IN $\text{AdS}_3(\text{BTZ BLACK HOLE}) \times$

$$\mathbf{S}^3 \times \mathbf{T}^4$$

A. Potential analysis

Here we consider the geometry of an exact $\text{AdS}_3(\text{AdS}_3 \times \mathbf{S}^3 \times \mathbf{T}^4)$ as the other solution to the type IIB string theory [9]. This corresponds to the non-dilatonic solution. This geometry can be led the BTZ black hole spacetime as a whole by the periodic identification. A ten-dimensional minimally coupled scalar satisfies

$$\square_{10} \Psi = 0. \quad (60)$$

Ψ can be decomposed into

$$\Psi = e^{-i\omega t} e^{im\varphi} e^{iK_i x^i} Y_l(\theta_1, \theta_2, \theta_3) \psi(\rho). \quad (61)$$

Then Eq.(60) leads to

$$\nabla_{\text{BTZ}}^2 \psi(\rho) + \frac{\mu}{R^2} \psi(\rho) = 0 \quad (62)$$

with $\mu = -l(l+2) - K^2 r_5^2$. The $\mu = -8(l=2)$ case contains both the dilaton(Φ) and a free scalar(ψ) with $l=2$. The $\mu = -3(l=1)$ case corresponds to an intermediate scalar (η) and $\mu = 1$ leads to tachyon(T). Here the BTZ black hole spacetime is given by [10]

$$ds_{\text{BTZ}}^2 = -f^2 dt^2 + \rho^2 \left(d\varphi - \frac{J}{2\rho^2} dt \right)^2 + f^{-2} d\rho^2 \quad (63)$$

with $f^2 = \rho^2/R^2 - M + J^2/4\rho^2 = (\rho^2 - \rho_+^2)(\rho^2 - \rho_-^2)/\rho^2 R^2$. The mass, angular momentum, angular velocity at horizon and area of horizon are

$$M = (\rho_+^2 + \rho_-^2)/R^2, \quad J = 2\rho_+ \rho_-/R, \quad \Omega_H = \frac{J}{2\rho_+^2}, \quad \mathcal{A}_H^{\text{BTZ}} = 2\pi\rho_+. \quad (64)$$

Further one finds the relation between the BTZ and a 5D black holes as

$$T_H^{\text{BTZ}} = (\rho_+^2 - \rho_-^2)/2\pi R^2 \rho_- = T_H, \quad \frac{1}{T_{L/R}^{\text{BTZ}}} = \frac{1}{T_H^{\text{BTZ}}} \left(1 \pm \frac{\rho_+}{\rho_-} \right) = \frac{1}{T_{L/R}}. \quad (65)$$

Its s-wave equation with $m=0$ takes the form

$$\left[f^2 \partial_\rho^2 + \left\{ \frac{1}{\rho} \partial_\rho (\rho f^2) \right\} \partial_\rho + \frac{\omega^2}{f^2} + \frac{\mu}{R^2} \right] \psi(\rho) = 0. \quad (66)$$

Defining $\psi(\rho) = \tilde{\psi}/\sqrt{\rho}$ and then (66) takes the form

$$f^2 \tilde{\psi}'' + (f^2)' \tilde{\psi}' + \left[\frac{f^2}{4\rho^2} - \frac{(f^2)'}{2\rho} + \frac{\omega^2}{f^2} + \frac{\mu}{R^2} \right] \tilde{\psi} = 0, \quad (67)$$

where the prime(') denotes the differentiation with respect to ρ . In order to obtain the Schrödinger-type equation, we introduce the tortoise coordinate ρ^* as [17]

$$\rho^* = \int \frac{d\rho}{f^2} = \frac{R^2}{2(\rho_+^2 - \rho_-^2)} \left[\rho_+ \ln \left(\frac{\rho - \rho_+}{\rho + \rho_-} \right) - \rho_- \ln \left(\frac{\rho - \rho_-}{\rho + \rho_+} \right) \right]. \quad (68)$$

We note that $\rho_+ \leq \rho \leq \infty$, while $-\infty \leq \rho^* \leq 0$ for outside horizon. On the other hand, in a 5D black hole one finds that $r_0 \leq r \leq \infty$ is mapped into $-\infty \leq r^* \leq \infty$. Then (67) leads to

$$\frac{d^2\tilde{\psi}}{d\rho^{*2}} + (\omega^2 - V_\mu) \tilde{\psi} = 0, \quad (69)$$

where the potential is given by

$$V_\mu(\rho) = f^2 \left[-\frac{f^2}{4\rho^2} + \frac{(f^2)'}{2\rho} - \frac{\mu}{R^2} \right]. \quad (70)$$

Four potential graphs ($V_T, V_\psi^{l=0}, V_\eta, V_\psi^{l=2}$) in an exact AdS_3 background are shown in Fig.4. The parameters are chosen as $\rho_+ = 0.01, \rho_- = 0.001, R = 0.3$. These are all monotonically increasing functions with ρ , contrast to V_N for an AdS_3 bubble. This shows a peculiar property of AdS_3 , which in this spacetime the asymptotic states cannot be defined. But the tachyon potential (V_T) is a monotonically decreasing function. This field couples to the minimal weight primary operator with $(1/2, 1/2)$ [18]. Also this satisfies both the stability condition for AdS_3 and the Dirichlet boundary condition [19], which are clearly related to a shape of its potential.




FIG. 4. Four potential graphs ($V_T, V_\psi^{l=0}, V_\eta, V_\psi^{l=2}$) as functions of ρ for an exact AdS_3 .

B. Asymptotically AdS₃-behavior and non-normalizable modes

In the near-horizon($\rho \rightarrow \rho_+$, $V_\mu \rightarrow 0$), (69) reduces to

$$\frac{d^2 \tilde{\psi}_{\text{NH}}}{d\rho^{*2}} + \omega^2 \tilde{\psi}_{\text{NH}} = 0 \quad (71)$$

which leads to the plane-wave solution

$$\tilde{\psi}_{\text{EH}} = e^{-i\omega\rho^*} + R_{\psi}^{\text{in}}(\omega)e^{i\omega\rho^*}. \quad (72)$$

Here the first term is an outgoing mode($\rightarrow$) and the second is an incoming mode($\leftarrow$). Now let us discuss the asymptotically AdS₃-behavior. Near the timelike boundary($\rho \rightarrow \infty$, $\rho^* \rightarrow 0$), one finds

$$\frac{d^2 \tilde{\psi}_\infty}{d\rho^{*2}} - \left[\frac{3}{4} - \mu \right] \frac{\rho^2}{R^4} \tilde{\psi}_\infty = 0. \quad (73)$$

Here we introduce the relation between ρ and ρ^*

$$\frac{\rho_+}{\rho(\rho^*)} = \frac{Y}{1 - \sigma^2} \sum_{n=0}^{\infty} a_n(\sigma) Y^{2n}, \quad Y = \tanh \lambda \rho^*, \quad \sigma = \frac{\rho_-}{\rho_+}, \quad \lambda = \frac{\rho_+}{R^2} (1 - \sigma^2), \quad a_0 = 1. \quad (74)$$

If σ^2 is very small and $\rho^* \rightarrow 0$, one finds

$$\rho = \frac{\rho_+}{Y} (1 - \sigma^2) = \rho_+ (1 - \sigma^2) \coth \lambda \rho^* \simeq \frac{R^2}{\rho^*}. \quad (75)$$

Using (75), (73) leads to

$$\frac{d^2 \tilde{\psi}_\infty}{d\rho^{*2}} - \left[\frac{3}{4} - \mu \right] \frac{\tilde{\psi}_\infty}{\rho^{*2}} = 0. \quad (76)$$

Its solution takes the form

$$\tilde{\psi}_\infty = \rho^{*\frac{1 \pm 2\sqrt{1-\mu}}{2}}. \quad (77)$$

Finally we have

$$\psi_\infty(\rho^*) = \frac{\tilde{\psi}_\infty}{\sqrt{\rho}} \propto \rho^{*(1 \pm \sqrt{1-\mu})}. \quad (78)$$

For the s-wave free scalar($\mu = 0$), ψ_∞ takes the form(ρ^{*2} , constant) and for the dilaton field($\mu = -8$), one finds $(\rho^{*4}, 1/\rho^{*2})$. We find $(\rho^{*3/2}, \rho^{*1/2})$ for a conformally coupled scalar($\mu = 3/4$) [17]. For the tachyon($\mu = 1$), one has (ρ^*, ρ^*) and for an intermediate scalar($\mu = -3$), one finds $(\rho^{*3}, 1/\rho^*)$. Instead of the plane-wave from, here one finds the power-law behaviors of ρ^* [20]. The non-normalizable modes are found as $1/\rho^{*2}$ for the dilaton and $1/\rho^*$ for an intermediate scalar, which diverge as ρ^* approaches the timelike boundary($\rho^* \rightarrow 0$). The positive powers of ρ^* all belong to be the normalizable modes, which converge at spatial infinity. We note that the tachyon takes only the normalizable modes [19]. This can be easily conjectured from its shape of potential V_T . The s-wave free scalar takes a constant-behavior at infinity. This makes an difficulty in dividing ψ_∞ into an ingoing and outgoing modes at spatial infinity [21]. Actually it is impossible to define an ingoing wave and an outgoing wave at spatial infinity of an exact AdS_3 . Instead in the asymptotically AdS_3 space it contains the normalizable as well as the non-normalizable modes. The latter will play an important role in calculation of the absorption coefficient $\mathcal{A}_\psi^{\text{in}}$.

C. Scattering from an exact AdS_3

First we note that ρ^* covers only the left-hand side($-\infty \leq \rho^* \leq 0$) of the whole space. In the case of where the black hole geometry is asymptotically flat as a 5D black hole, the tortoise coordinate r^* goes from $-\infty$ to ∞ . Hence this is similar to the infinite string problem in which the initial data propagates towards left and right indefinitely [17]. The initial data no longer enjoys this privilege when the background is asymptotically AdS_3 because the tortoise coordinate ρ^* goes from $-\infty$ to 0 only. One may consider this as the semi-infinite string problem(or a finite cavity with reflecting walls in AdS_5 [22]). Then the Dirichlet or Neumann boundary condition at spatial infinity($\rho^* = 0$) is required to formulate the problem appropriately. However, we take a different point of view to attack an asymptotically AdS_3 problem. This is based on the observation of the shape of potential

and the global structure of an exact AdS_3 [10]. We first construct the potential $V_\mu(\rho^*)$ by replacing ρ in (70) with $\rho \simeq \rho_+ \coth(\rho_+ \rho^*/R^2)$. The potentials in Fig.5 look like the half of the AdS_3 -bubble potentials. Especially, we observe that the free field potentials $V_\psi^{l=0}(\rho^*)$ and $V_\psi^{l=2}(\rho^*)$ in Fig.5 take nearly the same form as in $V_\phi^{l=0}(r^*)$ and $(V_\phi^{l=2}(r^*), V_\nu(r^*), V_\lambda(r^*))$ in the region of $-\infty \leq r^* \leq 0$ in Fig.3. Furthermore the Penrose diagram of an exact $\text{AdS}_3(-\infty \leq \rho^* \leq 0)$ is an half of the would-be whole diagram in $-\infty \leq \rho^* \leq \infty$ [10].




FIG. 5. Four potential graphs ($V_T, V_\psi^{l=0}, V_\eta, V_\psi^{l=2}$) as functions of ρ^* for an exact AdS_3 .

In this work we assume that the potential of an exact AdS_3 is the left-hand side one of an AdS_3 bubble. The important thing is to calculate the absorption cross section in the background of an exact AdS_3 . Considering the $\{\text{in}\}_\psi$ -state picture, it is not hard to calculate the absorption coefficient $\mathcal{A}_\psi^{\text{in}}$. Although the $\{\text{out}\}_\psi$ -state cannot be defined, we can derive $\mathcal{A}_\psi^{\text{in}}$ from the backscattering of a test field ψ off $V_\psi(\rho^*)$. If we choose the boundary condition appropriately, the of potential Fig.5 is enough to calculate the absorption coefficient, regardless of the would-be right-hand side ($0 \leq \rho^* \leq \infty$). In this backscattering process, the key point is to use an appropriate matching procedure between a near AdS_3 and an asymptotically

AdS₃. We remind the reader that $\rho^* = 0$ is a timelike boundary and thus information enter or exit from it. This is an exact middle point if one assumes a whole space of $-\infty \leq \rho^* \leq \infty$. Thus requiring the conventional boundary condition may lead to a wrong result in deriving the absorption coefficient. Instead of the Dirichlet condition of $\psi|_{\rho^*=0} = 0$, one may use the non-normalizable modes. The non-normalizable mode is a divergent quantity at $\rho^* = 0$ but its flux is finite at $\rho^* = 0$. Also it corresponds to specifying another boundary condition at spatial infinity. On the other hand, if one use the normalizable modes which satisfy the Dirichlet boundary condition, one may not succeed in obtaining the absorption coefficient in an exact AdS₃.

D. AdS₃-AdS₃ matching procedure and absorption cross section

In order to calculate the semiclassical absorption cross section, we have to solve the exact differential equation Eq.(62) with an appropriate boundary condition. Since it is difficult to solve Eq.(62) directly, one has to use the matching procedure between the near-horizon($\rho \sim \rho_+$) AdS₃ and the far-region($\rho \rightarrow \infty$) AdS₃. Here the matching point resides on $-\infty < \rho^* < 0$ in Fig.5.

In the far region Eq.(62) becomes

$$\psi''_{\infty} + \frac{3}{x}\psi'_{\infty} + \frac{\mu}{x^2}\psi_{\infty} = 0 \quad (79)$$

with a dimensionless variable $x = \rho/R$. One easily finds the far-region solution

$$\psi_{\infty}(x) = \left[\tilde{\alpha}x^{-1+\sqrt{1-\mu}} + \tilde{\beta}x^{-1-\sqrt{1-\mu}} \right] \quad (80)$$

with two unknown constants $\tilde{\alpha}, \tilde{\beta}$. The first term is a divergent quantity at $\rho = \infty$ but behaves well in the interior region. This corresponds to the non-normalizable modes and is coupled to the boundary operator $\mathcal{O}$ at infinity. The second one is the normalizable mode and propagates in the bulk. This can be used to construct two, three and four-point functions. Although one cannot define an ingoing flux at infinity of AdS₃, one can calculate the total flux. The flux at spatial infinity is given by

$$\mathcal{F}(\infty) = -2\pi\sqrt{1-\mu}\left|\tilde{\alpha} - i\tilde{\beta}\right|^2. \quad (81)$$

We note here that for the tachyon with $\mu = 1$, $\mathcal{F}^T(\infty) = 0$. This is because the tachyon takes only the normalizable modes which are to be zero at infinity. Thus we exclude it from our analysis.

In order to obtain the near-horizon behavior, we introduce the new variable $z = (\rho^2 - \rho_+^2)/(\rho^2 - \rho_-^2) = (x^2 - x_+^2)/(x^2 - x_-^2)$. Then Eq.(62) leads to

$$z(1-z)\frac{d^2\psi}{dz^2} + (1-z)\frac{d\psi}{dz} + \left(\frac{A_1}{z} + \frac{\mu/4}{1-z} - B_1\right)\psi = 0, \quad (82)$$

where

$$A_1 = \left(\frac{\omega - m\Omega_H}{4\pi T_H}\right)^2, \quad B_1 = -\frac{\rho_-^2}{\rho_+^2} \left(\frac{\omega - m\Omega_H \rho_+^2/\rho_-^2}{4\pi T_H^{\text{BTZ}}}\right)^2. \quad (83)$$

In the case of the s-wave propagation with $m = 0$, the near-horizon AdS_3 equation (82) leads exactly to the near-horizon equation (32) of a 5D black hole. Explicitly, the relationship between these is given by

$$A_1 \rightarrow Q, \quad \frac{\mu}{4} \rightarrow E, \quad B_1 \rightarrow C. \quad (84)$$

In this sense, (32) is called an AdS_3 bubble. The ingoing wave is given by the hypergeometric function

$$\psi(z) = C_1 z^{-i\sqrt{A_1}} (1-z)^{(1-\sqrt{1-\mu})} F(a, b, c; z), \quad (85)$$

where

$$\sqrt{1-\mu} \simeq l+1, \quad (86)$$

$$a = \frac{1-\sqrt{1-\mu}}{2} - i\sqrt{A_1} + i\sqrt{B_1} \simeq -\frac{l}{2} - i\frac{\omega}{4\pi T_R^{\text{BTZ}}}, \quad (87)$$

$$b = \frac{1-\sqrt{1-\mu}}{2} - i\sqrt{A_1} - i\sqrt{B_1} \simeq -\frac{l}{2} - i\frac{\omega}{4\pi T_L^{\text{BTZ}}}, \quad (88)$$

$$c = 1 - 2i\sqrt{A_1} = 1 - i\frac{\omega}{2\pi T_H^{\text{BTZ}}}. \quad (89)$$

Here $\simeq$ means $K^2 r_5^2 \simeq 0$. The corresponding flux is

$$\mathcal{F}(0) = -8\pi\sqrt{A_1}(x_+^2 - x_-^2) |C_1|^2 \quad (90)$$

with $x_+^2 - x_-^2 = (r_0/R)^2 \ll 1$. The absorption coefficient will be taken as

$$\mathcal{A}_\psi^{\text{in}} = \frac{\mathcal{F}(0)}{\mathcal{F}(\infty)} = \frac{4\sqrt{A_1}(x_+^2 - x_-^2)}{\sqrt{1-\mu}} \frac{|C_1|^2}{|\tilde{\alpha} - i\tilde{\beta}|^2}. \quad (91)$$

In order to obtain $\tilde{\alpha}$ and $\tilde{\beta}$, we use the matching procedure. It is important to remember that the present spacetime is an exact AdS_3 . Thus we have to match the near- AdS_3 with asymptotically AdS_3 to find an absorption coefficient. We know the far-region behavior of (85). This can be found from the $z \rightarrow 1 - z$ for the hypergeometric function

$$\psi_{n \rightarrow f}(x) = \left[C_1 E_1 (x_+^2 - x_-^2)^{(1-\sqrt{1-\mu})/2} x^{-1+\sqrt{1-\mu}} + C_1 E_2 (x_+^2 - x_-^2)^{(1+\sqrt{1-\mu})/2} x^{-1-\sqrt{1-\mu}} \right], \quad (92)$$

where

$$E_1 = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, E_2 = \frac{\Gamma(c)\Gamma(-c+a+b)}{\Gamma(a)\Gamma(b)}. \quad (93)$$

Matching (80) with (92) in the far-region ($x \gg 1$) leads to

$$\tilde{\alpha} = C_1 E_1 \left(\frac{r_0}{R} \right)^{1-\sqrt{1-\mu}}, \tilde{\beta} = C_1 E_2 \left(\frac{r_0}{R} \right)^{1+\sqrt{1-\mu}}. \quad (94)$$

Considering $R \gg r_0$, one finds $\tilde{\alpha} \gg \tilde{\beta}$ even for $l = 0$ case. Hence we can neglect $\tilde{\beta}$ in favor of $\tilde{\alpha}$. This amounts to taking the flux of the non-normalizable modes. The key point is an AdS_3 - AdS_3 matching in this backscattering process. Then the absorption coefficient is approximately given by

$$\mathcal{A}_\psi \simeq \frac{4\sqrt{A_1}(x_+^2 - x_-^2)}{\sqrt{1-\mu}} \left(\frac{r_0}{R} \right)^{2(\sqrt{1-\mu}-1)} \frac{1}{|E_1|^2}. \quad (95)$$

The absorption cross section for $\text{AdS}_3 \times \text{S}^3$ with $m = 0$ leads to

$$\sigma_{\text{AdS}}^\psi = \frac{\mathcal{A}_\psi}{\omega} \simeq \frac{\tilde{\mathcal{A}}_H^{6\text{D}}}{l!(l+1)!} \left(\frac{r_0}{R} \right)^{2l} \left| \frac{\Gamma(\frac{l+2}{2} - i\frac{\omega}{4\pi T_L^{\text{BTZ}}}) \Gamma(\frac{l+2}{2} - i\frac{\omega}{4\pi T_R^{\text{BTZ}}})}{\Gamma(l - i\frac{\omega}{2\pi T_H^{\text{BTZ}}})} \right|^2 \quad (96)$$

with $\tilde{\mathcal{A}}_H^{6\text{D}} = \mathcal{A}_H^{\text{BTZ}} \times 2\pi^2 R^3$.

In the low energy limit $\omega \rightarrow 0$, it turns out that the 6D cross sections for an exact AdS -theory take the same form as (55) and (57)

$$\sigma_{\text{AdS}}^{\psi_0} = \tilde{\mathcal{A}}_H^{6\text{D}}, \quad (97)$$

$$\sigma_{\text{AdS}}^{\psi_2} = \sigma_{\text{AdS}}^{\Phi} = \frac{1}{3} \frac{\tilde{\mathcal{A}}_H^{6\text{D}}}{4} \left(\frac{r_0}{R} \right)^4. \quad (98)$$

This 6D result is derived from $\text{AdS}_3 \times \text{S}^3$. In order for this to match with the cross section of a 5D black hole, it needs to introduce a compactified circle(S^1) in $\text{M}_5 \times \text{S}^1 \times \text{T}^4$. In this case one finds $\tilde{\mathcal{A}}_H^{6\text{D}} = \mathcal{A}_H^{5\text{D}} \times 2\pi R$ with a radius of $\text{S}^1(R)$. We note that ϕ_2 , $\nu(= \Phi)$ and λ give us slightly different cross sections in an AdS_3 bubble, whereas these(ψ_2, Φ) do not make any distinction in an exact AdS-theory.

IV. DISCUSSIONS

It seems that the S-matrix cannot be extracted from the anti-de Sitter space even in a limit [22,23]. This is based on the fact that in an exact AdS_3 the asymptotic states cannot be defined, due to the timelike boundary and the periodicity of geodesics. However, the authors in [8] showed that the correlation functions of the dual CFT_4 to AdS_5 are considered as the bulk S-matrices. The vacuum correlators $\langle \mathcal{O}(x_1) \mathcal{O}(x_2) \cdots \mathcal{O}(x_n) \rangle_{\text{CFT}_4}$ of the CFT_4 are expressed as truncated n-point functions convolved against the non-normalizable modes. These can be interpreted as an S-matrix for an exact AdS_5 arising from a limit of scattering from an AdS_5 bubble in asymptotically flat space.

In this work, we show that the S-matrix of an AdS_3 bubble can be derived from an exact AdS_3 in the dilute gas and low energy limits. We confirm this from the calculation of the absorption cross section. This originates from the fact that the near-horizon equations for an AdS_3 bubble (32) and an exact AdS_3 (82) are the same form, but they have different boundary conditions at infinity. In the AdS_3 bubble-calculation, one uses the AdS_3 -AFS matching to obtain the absorption coefficient $\mathcal{A}_N^{\text{out}}$. On the other hand, in the exact AdS_3 -calculation we use both the $\{\text{in}\}$ -state picture and the non-normalizable mode to obtain the absorption coefficient $\mathcal{A}_\psi^{\text{in}}$. This amounts to taking the AdS_3 - AdS_3 matching. The s-wave greybody factor of a free scalar of an AdS_3 bubble has exactly the same form as that of an

exact AdS_3 . For the dilaton we find the same form of cross section $\sigma = c\tilde{\mathcal{A}}_H^{6\text{D}}(r_0/R)^4$ but with $c = 1/4$ for an AdS_3 bubble and $c = 1/12$ for an exact AdS_3 .

Let us compare our results with the others. The general formular for the gerybody factor is derived from the vacuum two-point function $\langle \mathcal{O}(x)\mathcal{O}(0) \rangle_{\text{CFT}_2}$ of a boundary operator $\mathcal{O}$ in the effective string [24] and boundary CFT_2 approaches [25]. These give us the same result for a free scalar but for the dilaton, $c = 1/4$ as in an AdS_3 bubble. Consequently, the two-point correlator provides us the greybody factor in the dilute gas and low energy limits. This quantity takes exactly the same form in the CFT_2 and AdS_3 bubble approaches. Further, in the exact AdS_3 approach one finds the same form of the greybody factor. This means that the S-matrix can be derived from an exact AdS_3 space. It is obvious that the conformal limit of the gauge theory(CFT_2) corresponds with scattering from an exact AdS_3 . Finally we present here a scattering picture in an exact AdS_3 and compare it with the scattering of an AdS_3 bubble in AFS.

ACKNOWLEDGMENTS

This work was supported in part by the Basic Science Research Institute Program, Ministry of Education, Project NOs. BSRI-98-2413 and BSRI-98-2441.

REFERENCES

- [1] A. Strominger and C. Vafa, Phys. Lett. **B379**, 99 (1996), hep-th/9601029; C. Callan and J. Maldacena, Nucl. Phys. **B472**, 591 (1996), hep-th/9602043; G. Horowitz and A. Strominger, Phys. Rev. Lett. **77**, 2368 (1996), hep-th/9602051.
- [2] A. Dhar, G. Mandal, and S. Wadia, Phys. Lett. **B388**, 51 (1996), hep-th/9605234.
- [3] S. Das, G. Gibbons, and S. Mathur, Phys. Rev. Lett. **78**, 417 (1997).
- [4] C. Callan, S. Gubser, I. Klebanov, and A. Tseytlin, Nucl. Phys. **B489**, 65 (1997), hep-th/9610172; I.R. Klebanov and M. Krasnitz, Phys. Rev. **D55**, R3250 (1997).
- [5] M. Krasnitz and I. Klebanov, Phys. Rev. **D56**, 2173 (1997), hep-th/9703216; M. Taylor-Robinson, hep-th/9704172; H. W. Lee, Y. S. Myung, J. Y. Kim, Phys. Rev. **D58**, 104006(1998), hep-th/9708099.
- [6] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231(1998), hep-th/9711200; S.S. Gubser, I.R. Klebanov and A.M. Polyakov, Phys. Lett. **B428**, 105(1998), hep-th/9802109; E. Witten, Adv. Theor. Math. Phys. **2**, 253(1998), hep-th/9802150.
- [7] S. Hyun, hep-th/9704005 ; K. Sfetsos and K. Skenderis, Nucl. Phys. **B517**, 179 (1998), hep-th/9711138; S. de Alwis, Phys. Lett. **B435**, 31 (1998), hep-th/9804019.
- [8] V. Balasubramanian, S. Giddings, and A. Lawrence, JHEP **9903**, 001 (1999), hep-th/9902052.
- [9] H. W. Lee and Y. S. Myung, Phys. Rev. **D58**, 104013 (1998), hep-th/9804095; H.W. Lee, N.J. Kim, and Y.S. Myung, hep-th/9805050.
- [10] M. Banados, M. Henneaux, C. Teitelboim, and A. Zanelli, Phys. Rev. **D48**, 1506(1993).
- [11] J. Maldacena and A. Strominger, Phys. Rev. **D56**, 4975 (1997), hep-th/9702015;
- [12] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1403 (1957); C. V. Vishveshwara, Phys.

- Rev. **D1**, 2870 (1970); F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970); O. J. Kwon, Y. D. Kim, Y. S. Myung, B. H. Cho and Y. J. Park, Phys. Rev. **D34**, 333 (1986).
- [13] R. Dijkgraaf, H. Verlinde and E. Verlinde, Nucl. Phys. **B371**, 269(1992); K. Sfetsos, Nucl. Phys. **B389**, 424(1993); J. Y. Kim, H. W. Lee and Y. S. Myung, Phys. Rev. **D50**, 3942 (1994) ; Y. S. Myung, Phys. Lett. **B362**, 46 (1995).
- [14] S. Chandrasekhar, *The Mathematical Theory of Black Hole* (Oxford Univ. Press, New York, 1983).
- [15] S.W. Hawking, Comm. Math. Phys. **43**, 199(1975); N. Birrell and P. Davies, *Quantum fields in curved space* (Cambridge University Press, 1982, N.Y.).
- [16] S.D. Mathur, Nucl. Phys. **B514**, 204(1998), hep-th/9704156.
- [17] J.S.F. Chan and R.B. Mann, Phys. Rev. **D55**, 7546(1997).
- [18] J. Maldacena and A. Strominger, JHEP **9812**, 005(1998), hep-th/9804085.
- [19] H.W. Lee, N.J. Kim, and Y.S. Myung, Phys. Lett. **B441**, 83(1998), hep-th/9803227.
- [20] V. Balasubramanian, P. Krauss, and A. Lawrence, Phys. Rev. **D59**, 046003 (1999), hep-th/9805171.
- [21] D. Birmingham, I. Sachs and S. Sen, Phys. Lett. **B413**, 281(1997); H.W. Lee, N.J. Kim and Y.S. Myung, Phys. Rev. **D58**, 084022(1998), hep-th/9803080.
- [22] L. Susskind, hep-th/9901079.
- [23] J. Polchinski, hep-th/9901076.
- [24] S.S. Gubser, Phys. Rev. **D56**, 7854(1997).
- [25] E. Teo, Phys. Lett. **B436**, 269(1998).